

HOT MOLECULAR GAS IN THE GALACTIC CENTER

ROBESON M. HERRNSTEIN¹ AND PAUL T.P. HO¹*Accepted for publication in ApJ Letters*

ABSTRACT

Using the new 23 GHz receivers at the Very Large Array (VLA), we have detected NH₃(6,6) emission ($\nu = 25.056025$ GHz) from hot (> 150 K) molecular clouds in the central 10 pc of the Galaxy. This is the first successful detection of NH₃(6,6) with the VLA. The brightest emission comes from a region interior to the “circumnuclear disk” (CND), less than 1.5 pc in projected distance from Sgr A*. This region does not show molecular emission from lower energy transitions such as NH₃(1,1) and (2,2), HCN(1-0) and HCO⁺(1-0). Line ratios of NH₃(6,6) and (3,3) emission as well as NH₃(6,6) line widths have peak values within 1.5 pc of Sgr A*, indicating that the gas is physically close to the nucleus. NH₃(6,6) is also detected towards many features outside the CND observed in NH₃(1,1), (2,2), and (3,3). These features tend to lie along ridges of gas associated with Sgr A East or the massive “molecular ridge” that connects the “20 km s⁻¹” and “50 km s⁻¹” giant molecular clouds (GMCs).

Subject headings: Galaxy: center — ISM: clouds — ISM: molecules — radio lines: ISM

1. INTRODUCTION

At a distance of only 8.0 ± 0.5 kpc (Reid 1993), the Galactic center provides a unique opportunity to study in detail the environment around a supermassive black hole. It is now generally accepted that a black hole of $2.6 \times 10^6 M_{\odot}$ is located at the dynamical center of the Galaxy (Eckart & Genzel 1997; Ghez et al. 1998). In the radio, emission from just outside the black hole is observed as the strong (~ 1 Jy) source, Sgr A*. Sgr A* is surrounded by arcs of ionized gas (Sgr A West) that appear to be feeding the nucleus (Lo & Claussen 1983; Roberts & Goss 1993). These arcs are, in turn, surrounded by an apparent “ring” of molecular material called the circumnuclear disk (CND, Güsten et al. (1987)).

Sgr A West and the CND are located in front of or just inside the front edge of the expanding supernova remnant (SNR), Sgr A East (Pedlar et al. 1989). The expansion of Sgr A East appears to be moving large amounts of material away from the nucleus, forming ridges of material on all sides (Genzel et al. 1990; Ho et al. 1991; Serabyn, Lacy, & Achtermann 1992; Zylka et al. 1999; McGary, Coil, & Ho 2001, hereafter MCH). However, some material may move towards the nucleus after being disrupted by the passing front and three filamentary “streamers” possibly feeding the nucleus have been detected in NH₃ (Okumura et al. 1989; Ho et al. 1991; Coil & Ho 1999, 2000; MCH).

The metastable (J=K) NH₃(J,K) rotation inversion transitions at ~ 23 GHz have proven to be useful probes of dense (10^4 – 10^5 cm⁻³) molecular material near the Galactic center. They tend to have a low optical depth and a high excitation temperature at the Galactic center, making them almost impervious to absorption effects. Satellite hyperfine lines separated by 10–30 km s⁻¹ on either side of the main line enable a direct calculation of the optical depth of the NH₃ emission and line ratios of different transitions can be used to calculate the rotational temperature,

T_R , of the gas.

We recently observed NH₃(1,1), (2,2) and (3,3) emission from the central 10 pc of the Galaxy. An important result from these studies is the apparent increase in line width and T_R as gas approaches the nucleus (MCH; McGary & Ho in press). However, the reality of this effect has remained in doubt because the emission also becomes fainter near Sgr A*. We suspected that even the NH₃(3,3) line becomes less sensitive to the extreme environment near Sgr A* and observed the central 4' (10 pc) of the Galaxy in NH₃(6,6), at 412 K above ground, using the new 23 GHz receivers at the Very Large Array² (VLA) in order to detect the hottest molecular gas. These data represent the first successful observations of NH₃(6,6) with the VLA. NH₃(6,6) is detected in many of the features seen in lower NH₃ transitions, but the velocity integrated map is dominated by emission less than 1.5 pc (40'') in projected distance from the nucleus. Line ratios and extremely broad line widths (50–90 km s⁻¹) indicate this molecular gas is physically close to the supermassive black hole. The remaining features lie predominantly along the edge of Sgr A East and have line widths greater than 20 km s⁻¹ indicating interaction with the shell.

2. OBSERVATIONS

Observations of NH₃(6,6) were made with the VLA in the D north-C array on 2001 October 1 and November 16 and have an identical setup and spatial coverage to our previous NH₃(1,1), (2,2), and (3,3) observations so that comparisons between images are straightforward (MCH). A five-pointing mosaic centered on Sgr A* ($\alpha_{2000} = 17^h45^m40^s.0$, $\delta_{2000} = -29^\circ00'26''.6$) fully samples the central 4' (10 pc) of the Galaxy and a velocity coverage of ± 120 km s⁻¹ with a resolution of 9.3 km s⁻¹ includes most of the velocities observed near the nucleus (the features at -185 km s⁻¹ observed by Zhao, Goss,

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, rmccgary@cfa.harvard.edu, pho@cfa.harvard.edu

² The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

& Ho (1995) are not covered). The data have been calibrated and images made using the same method as MCH. The final beam size after application of a Gaussian taper (FWHM=10'') to the uv data is $12.0'' \times 9.2''$ with a PA of -1.52° .

Observations at 25 GHz have only recently become possible after the upgrade of the 23 GHz receivers at the VLA. At the time of our observations, 16 antennas were fitted with new receivers. At the low declination of Sgr A*, the new receivers have $\langle T_{sys} \rangle = 96$ K, which is comparable to the noise at 23 GHz with the old receivers. Data are weighted by $1/T_{sys}^2$ so that old receivers do not increase the noise. One hour of integration time was obtained for each pointing resulting in an rms noise per channel of $\sigma_{ch} = 8.7$ mJy beam $^{-1}$. For the velocity integrated image, the 1σ noise level is 0.23 Jy beam $^{-1}$ km s $^{-1}$ and is calculated in the same way as MCH assuming line emission typically appears in seven channels.

3. RESULTS

Figure 1 shows velocity integrated NH₃(6,6) emission in contours overlaid on a pseudo-color 6 cm continuum image (Yusef-Zadeh & Morris 1987). The point source Sgr A* ($\Delta\alpha = 0''$, $\Delta\delta = 0''$) is the brightest feature in the 6 cm image. The arcs of Sgr A West can be seen in yellow and the large expanding shell, Sgr A East, is seen in blue. The contours for NH₃(6,6) are in steps of 3σ . Figure 2 shows the velocity integrated map with labeled positions for twelve spectra shown to the right. The spectra have been Hanning smoothed to a resolution of ~ 20 km s $^{-1}$. Spectrum L shows the noise at a position with no velocity integrated emission. Note the presence of many separate features at the position of Sgr A* (spectrum A). These features are due to separate clouds, and not satellite hyperfine lines which are only ~ 30 km s $^{-1}$ from the main line. With the exception of the 70 km s $^{-1}$ feature, the emission is not coincident with the velocities of known clouds in the region, nor do we expect line-of-sight (LOS) absorption associated with this high T_R . For spectra B–K, the best fit to a Gaussian is overlaid and the line width (defined as the FWHM) in km s $^{-1}$ is labeled in the upper right-hand corner of each panel. The NH₃(6,6) line widths are high throughout the central 10 pc. The typical line width for cool molecular gas at the Galactic center is ~ 15 km s $^{-1}$ (Armstrong & Barrett 1985; MCH), but none of the NH₃(6,6) features has a line width of less than 20 km s $^{-1}$, despite our sensitivity to FWHM $\gtrsim 6$ km s $^{-1}$.

Finally, we show the line ratio of NH₃(6,6) to (3,3) emission in color in Figure 3. To match the resolution of the NH₃(3,3) image, the NH₃(6,6) image was remade using a Gaussian taper resulting in $15'' \times 13''$ resolution and $\sigma'_{ch} = 11$ mJy beam $^{-1}$, which is equal to the noise in the (3,3) image, σ_{33} (MCH). The velocity channel corresponding to peak NH₃(6,6) emission is used to measure both the (6,6) and (3,3) flux densities which are then used to calculate the line ratios. Only pixels with NH₃(6,6) flux density greater than $3\sigma'_{ch}$ are included. For those pixels where NH₃(3,3) emission is less than $3\sigma_{33}$, the ratio of (6,6) emission to $3\sigma_{33}$ is used as a lower limit for the line ratio. Contours of the velocity integrated NH₃(6,6) emission are overlaid for reference.

If NH₃(6,6) and (3,3) are in thermal equilibrium, the

line ratio gives an estimate of the rotational temperature of the gas. From Townes & Schalow (1975), T_R is related to the opacity of the NH₃(3,3) main line and the ratio of NH₃(6,6) and (3,3) antenna temperatures by

$$T_R = \frac{-287 \text{ K}}{\ln\left[\frac{-0.43}{\tau_m(3,3)} \ln\left(1 - \frac{\Delta T_A(6,6)}{\Delta T_A(3,3)} (1 - e^{-\tau_m(3,3)})\right)\right]} \quad (1)$$

if equal beam filling factors are assumed. The exponential dependence of T_R on the line ratio reduces the accuracy of the estimation for $T_R \gtrsim 1000$ K. This is a significant improvement on NH₃(2,2) to (1,1) line ratios which only give accurate estimations for $T_R \lesssim 50$ K. The rotational temperature is thought to be a good indicator of kinetic temperature, T_K , and we assume T_R gives a lower limit for T_K (Martin, Ho, & Ruf 1982).

Line ratios greater than 2.3 are seen in Figure 3, but they give $T_R < 0$. These large line ratios may be the result of a larger filling factor for NH₃(6,6), or the dynamic range of the NH₃(3,3) data may be limited by nearby bright emission (MCH). It is unlikely that the gas is out of thermal equilibrium because the equilibration time is only $\sim 10^3$ s. Line widths are quite large in the region ($50 - 80$ km s $^{-1}$) making it unlikely for the NH₃(6,6) population to be inverted. The large line widths also make absorption of NH₃(3,3) by an un-associated, cool foreground cloud unlikely. If the NH₃(6,6) originates in a radiatively heated cloud, then it is possible that the NH₃(3,3) is absorbed by cooler material in the same cloud that has been shielded from the radiation. The NH₃(6,6) would be unaffected by absorption because the cooler gas would contain almost no NH₃(6,6). For clouds near the nucleus, the radiation involved would most likely originate in the central stellar clusters and those clouds with line ratios > 2.3 would be located in front of the nucleus along the LOS.

3.1. The central 1.5 parsecs

The NH₃(6,6) velocity integrated map is dominated by emission less than 1.5 pc ($40''$) from Sgr A*, interior to the CND (spectra B, C, & D). Indications of molecular gas inside the CND have also been observed in NH₃(3,3), but the emission was much fainter than surrounding features (MCH). Observations of $63 \mu\text{m}$ emission from [OI] with a resolution of $22''$ indicate a central concentration of neutral gas interior to the CND, possibly associated with the arcs of Sgr A West (Jackson et al. 1993). The distribution of the NH₃(6,6) does not mimic the shape of Sgr A West, and the line profiles do not show the high velocities associated with the ionized arcs and [OI] emission (Jackson et al. 1993; Roberts & Goss 1993).

The NH₃(6,6) emission appears to be kinematically independent of material in the CND. To the north of Sgr A* (spectra B and C), the gas has a velocity of ~ 0 km s $^{-1}$, but HCN(1-0) and HCO⁺(1-0) emission from nearby parts of the CND have velocities of $50 - 100$ km s $^{-1}$ (Wright et al. 2001). To the southeast of Sgr A* (spectrum D) emission is coincident with a gap in the CND and possible origins for this cloud are discussed at the end of this section. Spectrum E has a velocity of -20 km s $^{-1}$ in NH₃(6,6) compared to -5 km s $^{-1}$ in NH₃(3,3). HCN(1-0) and HCO⁺(1-0) spectra at this position are affected by absorption and central velocities cannot be determined, but emission from the nearby southern lobe of the CND is centered at -100 km s $^{-1}$ (Wright et al. 2001). The NH₃(6,6)

line has a relatively small line width (48 km s^{-1}) and low line ratio (0.3) compared to spectra B, C, and D and we assume this feature is not located in the central parsec.

The large line widths of $75\text{--}85 \text{ km s}^{-1}$ in the central 1.5 pc (spectra B, C, and D) indicate that this gas is physically close to the nucleus. The line profiles are similar to those from the CND for other molecules (e.g. Wright et al. (2001)) and are well-fitted by Gaussians. The smooth profiles make turbulence an unlikely broadening mechanism. Observations of the expansion of SNRs into molecular material show that the resulting line profiles are often best fit by a combination of dissociative J-type and non-dissociative C-type shocks (Wang & Scoville 1992; van Dishoeck, Jansen, & Phillips 1993). If the material is traveling perpendicular to the LOS, the combination of the two shocks results in a symmetrically broadened profile (van Dishoeck, Jansen, & Phillips 1993). Thus an expanding shock front perpendicular to our LOS may explain the large observed line widths.

The $\text{NH}_3(6,6)$ to $(3,3)$ line ratios are very high for spectra B, C, and D and many of the features in the central 1.5 pc exceed the upper limit for the line ratio of 2.3. A direct calculation of T_R is not possible, but the trends in Figure 3 indicate that gas near the nucleus is hot and we assume a warm kinetic temperature of 500 K . For an approximate mass estimate, we assume an optical depth of 0.1 and $N_{\text{H}_2} = 10^8 N_{\text{NH}_3}$ (Serabyn & Güsten 1986) and find $N_{\text{H}_2} = 8 - 9 \times 10^{23} \text{ cm}^{-2}$ for B, C, and D, and a total gas mass in the central 2 pc on the order of $10^4 M_\odot$. This mass is similar to those calculated by (Coil & Ho 1999, 2000) for cooler molecular clouds in the central 10 pc . The estimated virial mass is $\sim 10^6 M_\odot$ for a cloud of radius $15''$ and line width 80 km s^{-1} (MacLaren, Richardson, & Wolfendale 1988). Thus, it is safe to assume that these clouds are not gravitationally bound.

In lower transitions of NH_3 , there is evidence for a connection of the “ 20 km s^{-1} ” GMC to the nucleus along the “southern streamer” ($\Delta\alpha = 30''$, $\Delta\delta = -100''$) (Ho et al. 1991; Coil & Ho 1999, 2000). The gas appears to become warmer and line widths broaden as the southern streamer approaches the nucleus. However, the emission disappears within 2 pc of the nucleus. We detect significant $\text{NH}_3(6,6) \sim 20''$ to the southeast of Sgr A*, coincident with the “gap” in the CND seen in $\text{HCN}(1-0)$ and $\text{HCO}^+(1-0)$ (Wright et al. 2001). This gas has a velocity of 40 km s^{-1} and appears to be kinematically connected to the southern streamer (spectrum I). The large line ratios and broadened line widths (up to 80 km s^{-1}) indicate that it may trace the continuation of the southern streamer towards the nucleus. The velocity is less than the escape velocity for a distance of 2 pc , but ambiguities in projected distance and velocity make it impossible to claim that it is definitely bound by the nucleus.

3.2. Features outside the CND

The “western streamer” is offset $1'$ to the west of Sgr A* along the edge of Sgr A East (coincident with spectra G & H). It is known to have high ratios of $\text{NH}_3(2,2)$ to $(1,1)$ emission (MCH) as well as large line widths (McGary & Ho in press). In $\text{NH}_3(6,6)$, line widths range between 30 and 50 km s^{-1} . The line ratios in the western streamer have an average value of 0.66 indicating $T_R \approx 230 \text{ K}$ if the gas is optically thin. The velocity gradient of $25 \text{ km s}^{-1} \text{ pc}^{-1}$

observed in NH_3 by MCH accounts for the line velocities of $+30 \text{ km s}^{-1}$ in spectrum G and -30 km s^{-1} in spectrum H. The gradient is easily explained by a long filament highly inclined to the the LOS and moving out with the expanding shell. The impact of the shell perpendicular to the LOS could again explain the broad line widths in this feature.

The molecular ridge that connects the “ 20 km s^{-1} ” GMC in the south to the “ 50 km s^{-1} ” GMC in the east (Ho et al. 1991; Dent et al. 1993; Coil & Ho 1999, 2000, MCH) is composed of the southern part of the southern streamer, SE1 ($70''$, $-100''$), and SE2 ($110''$, $-60''$) (spectra I, J, & K, respectively). Emission from the molecular ridge has a typical velocity of 35 km s^{-1} , line width of 25 km s^{-1} , and an average line ratio of 0.41 (170 K), making it relatively cool compared to other features in our map. However, the line widths are roughly 10 km s^{-1} larger than that associated with the nearby, cold GMCs. The concave southeastern edge of Sgr A East seen in the 6 cm image is thought to result from an impact with SNR G359.92–0.09 (centered at $\Delta\alpha = 120''$, $\Delta\delta = -180''$, (Coil & Ho 2000)) and it is possible that the shocks from this collision have produced increased temperatures and densities (Coil & Ho 2000). There is a lack of $\text{NH}_3(6,6)$ emission on the northeast edge of Sgr A East where it is impacting the massive, but relatively quiescent, “ 50 km s^{-1} ” GMC. The high density of this GMC may result in a faster cooling rate in this feature, limiting the production of $\text{NH}_3(6,6)$.

Within the outer edge of the Sgr A East shell is a well-known north-south ridge of continuum radio emission $\sim 50''$ to the east of Sgr A* (coincident with spectrum F), roughly along the eastern edge of Sgr A West (Pedlar et al. 1989). Although the northern end of the ridge is coincident with the eastern edge of the CND, $\text{NH}_3(3,3)$ observations indicate that the 50 km s^{-1} gas is most likely an extension of SE1 (MCH). The $\text{NH}_3(6,6)$ line width along this ridge ranges from $40\text{--}60 \text{ km s}^{-1}$ and has an average line ratio of 1.0 corresponding to $T_R = 340 \text{ K}$. The line ratio peaks in the south where there is a lack of emission from HCN (Wright et al. 2001). The similarity of the western streamer and this eastern ridge as well as the asymmetric line profile (spectrum F) indicates that it likely lies on the expanding front of Sgr A East (Pedlar et al. 1989).

4. SUMMARY

With 25 GHz observations now practical at the VLA, $\text{NH}_3(6,6)$ is an excellent tracer of hot, dense molecular material. Velocity integrated $\text{NH}_3(6,6)$ images are dominated by emission less than 1.5 pc in projected distance from Sgr A*. High line ratios and line widths indicate that this molecular material is likely close to the supermassive black hole and is interior to the CND. In addition, the hottest emission to the southeast of Sgr A* appears to connect the southern streamer to the nucleus. The remaining $\text{NH}_3(6,6)$ emission tends to lie along the ridges that surround the Sgr A East shell, tracing features that show a high line width in earlier $\text{NH}_3(1,1)$, $(2,2)$, and $(3,3)$ data.

The authors thank P. Sollins, T. Bergin, and the referee for helpful comments and D. Shepherd for help with the 25 GHz system. RSM is supported in part by a Harvard University Merit Fellowship.

REFERENCES

- Armstrong, J.T. & Barrett, A.H. 1985, *ApJS*, 57, 535
- Coil, A.L. & Ho, P.T.P. 1999, *ApJ*, 513, 752
- Coil, A.L. & Ho, P.T.P. 2000, *ApJ*, 533, 245
- Dent, W.R.F., Matthews, H.E., Wade, R. , & Duncan, W.D. 1993, *ApJ*, 410, 650
- Eckart, A. & Genzel, R. 1997, *MNRAS*, 284, 576
- Genzel, R., Stacey, G.J., Harris, A.I., Geis, N., Graf, U.U., Poglitsch, A. , & Stutzki, J. 1990, *ApJ*, 356, 160
- Ghez, A.M., Klein, B.L., Morris, M. , & Becklin, E.E. 1998, *ApJ*, 509, 678
- Güsten, R., Genzel, R., Wright, M.C.H., Jaffe, D.T., Stutzki, J. , & Harris, A.I. 1987, *ApJ*, 318, 124
- Herzberg, G. 1966, *Molecular Spectra and Molecular Structure* (New York: Van Nostrand Reinhold Co.)
- Ho, P.T.P., Ho, L.C., Szczepanski, J.C., Jackson, J.M., Armstrong, J.T. , & Barrett, A.H. 1991, *Nature*, 350, 309
- Ho, P.T.P. & Townes, C.H. 1983, *ARA&A*, 21, 239
- Jackson, J.M., Geis, N., Genzel, R., Harris, A.I., Madden, S., Poglitsch, A., Stacey, G.J., & Townes, C.H. 1993, *ApJ*, 402, 173
- Lo, K.Y. & Claussen, M.J. 1983, *Nature*, 306, 647
- MacLaran, I., Richardson, K.M., & Wolfendale, A.W. 1988, *ApJ*, 333, 821
- Marshall, J., Lasenby, A.N. , & Harris, A.I. 1995, *MNRAS*, 277, 594
- Martin, R.N., Ho, P.T.P., & Ruf, K. 1982, *Nature*, 296, 632
- McGary, R.S., Coil, A.L., & Ho, P.T.P. 2001, *ApJ*, 559, 326 (MCH)
- McGary, R.S. & Ho, P.T.P. 2002, *ApJ*, in press (astro-ph/0206077)
- Mezger, P.G., Zylka, R., Salter, C.J., Wink, J.E., Chini, R. Kreysa, E. , & Tuffs, R. 1989, *A&A*, 209, 337
- Okumura, S.K., et al. 1989, *ApJ*, 347, 240
- Pedlar, A., Anantharamaiah, K.R., Ekers, R.D., Goss, W.M., van Gorkom, J.H., Schwarz, U.J., & Zhao, J.-H. 1989, *ApJ*, 342, 769
- Reid, M.J. 1993, *ARA&A*, 31, 345
- Roberts, D.A. & Goss, W.M. 1993, *ApJS*, 86, 133
- Serabyn, E. & Güsten, R. 1986, *A&A*, 161, 334
- Serabyn, E., Lacy, J.H. , & Achermann, J.M. 1992, *ApJ*, 395, 166
- Townes, C.H. & Schalow, A.L. 1975, *Microwave Spectroscopy*, New York:Dover
- van Dishoeck, E.F., Jansen, D.J., & Phillips, T.G. 1993, *A&A*, 279, 541
- Wang, Z. & Scoville, N.Z. 1992, *ApJ*, 386, 158
- Wright, M.C.H.W., Coil, A.L., McGary, R.S., Ho, P.T.P. , & Harris, A.I. 2001, *ApJ*, 551, 254
- Yusef-Zadeh, F. & Morris, M. 1987, *ApJ*, 320, 545
- Zhao, J.-H., Goss, W.M., & Ho, P.T.P. 1995, *ApJ*, 450, 122
- Zylka, R., Güsten, R., Philipp, S., Ungerechts, H., Mezger, P.G. , & Duschl, W.J. 1999, in *ASP Conf. Ser.*, 186, *The Central Parsecs of the Galaxy*, ed. H. Falcke et al. (San Francisco: ASP), 415




FIG. 1.— $\text{NH}_3(6,6)$ velocity integrated emission in contours overlaid on 6 cm continuum emission in color (Yusef-Zadeh & Morris 1987). Contours are in steps of 3σ where $1\sigma = 0.23 \text{ Jy beam}^{-1} \text{ km s}^{-1}$.




FIG. 2.— *Left* : Position of spectra overlaid on velocity integrated $\text{NH}_3(6,6)$ emission. Sgr A* is located at (0,0). *Right* : $\text{NH}_3(6,6)$ spectra with Gaussian fits overlaid showing that the largest line widths are located close to the nucleus. Line widths in km s^{-1} are given for spectra B-K in the upper right-hand corner of each panel.




FIG. 3.— Ratio of main line $\text{NH}_3(6,6)$ and $(3,3)$ emission. The velocity channel for the main line is chosen using the $\text{NH}_3(6,6)$ image cube. Line ratios are calculated for every point with $(6,6)$ emission $> 3\sigma'_{ch}$. For pixels with faint $(3,3)$ emission, $3\sigma_{33}$ is used to estimate a lower limit for the line ratio. Sgr A* is located at $(0,0)$.